



STUDY OF PREVALENCE OF ASTHMA IN SCHOOL CHILDREN IN PUNE AND ITS RELATION WITH FAMILY HISTORY OF ASTHMA AND ALLERGIC DISORDERS

Sampada Tambolkar

Professor, Dr D.y. Patil Medical College, Hospital And Research Center& Dr. D.y.patil Vidyapeeth, Pimpri, Pune, Maharashtra, India

Ambrish Mishra

Associate Professor, Dr D.y. Patil Medical College, Hospital And Research Center& Dr. D.y.patil Vidyapeeth, Pimpri, Pune, Maharashtra, India

Manojkumar Patil*

Associate Professor, Dr D.y. Patil Medical College, Hospital And Research Center& Dr. D.y.patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.*Corresponding Author
Email id: manoj.patil@dpu.edu.in

Shradha Salunkhe

Associate Professor, Dr D.y. Patil Medical College, Hospital And Research Center& Dr. D.y.patil Vidyapeeth, Pimpri, Pune, Maharashtra, India

Sharad Agarkhedkar

Professor And Head Of Department, Dr D.y. Patil Medical College, Hospital And Research Center& Dr. D.y.patil Vidyapeeth, Pimpri, Pune, Maharashtra, India

ABSTRACT

Objective: To determine prevalence of asthma in Pune and its relation to family history of allergic disease. **Study**

Design: Cross-sectional study in two schools of Pune. **Method:** International Study of Asthma and Allergy in Childhood (ISAAC) based questionnaire was administered to 815 parents of students of 2 selected schools between ages of 5-15 years. Responses were analyzed and children with history suggestive of asthma were examined, investigated at a tertiary teaching hospital. **Results:** prevalence of asthma was 6.13%. Prevalence in males was 6.36% and in females was 8.05% (p= 0.639). Prevalence in English medium school was 7.36% as compared to Marathi medium school of 3.8%. Family history of asthma and allergic disease had strong association with asthma. PEFR at time of enrolment was decreased in many children irrespective of symptoms. **Conclusion:** Prevalence of asthma was 6.13%. Family history is strong determinant of asthma.

KEY WORDS : Asthma Prevalence, Family History, PEFR Correlation.

INTRODUCTION

Asthma is the most common chronic disease of childhood. It is characterized by airway hyper responsiveness and airway inflammation. Asthma is more common in children (8.6 %) than adults (7.4 %). (1) .Family history of asthma, atopic dermatitis, allergic rhinitis, food or inhalant allergy is common risk factors for persistent asthma. (1). Atopy is the single largest risk factor for development of asthma. Nearly 80% of atopic individuals develop asthma, allergic rhinitis, eczema or urticarial. (2). Atopy is associated with increased risk of persistent and severe asthma. Allergen provocation can induce bronchial inflammation. Children who developed early sensitization to food or aero antigen and had eczema are at high risk for developing persistent asthma. (3)Asthma is a global health care problem, with increasing prevalence being observed over last few decades. (1). Prevalence of asthma in developing countries is significantly different from developed countries. In India, prevalence varies widely due to the vast size of the country, variable population density, variable climate, variable pollution levels and variable life style. (4)ISAAC, largest study determining prevalence of asthma had showed prevalence of asthma in Pune of 2.9% in 2003 (5). In recent study of 2012, study conducted in Pune suggested an increase in prevalence of asthma from 2.9% to 6.7%. (4) This study defined asthma from history provided in questionnaire and no clinical examination was done to confirm the diagnosis. To substantiate this observation after confirming the diagnosis with examination, this study was planned. PEFR was performed as a marker of lung function.

METHODOLOGY

It was a cross sectional study done in 815 children (5-15 yrs.), from two urban schools (English medium and Marathi medium each). Questionnaire - adopted from International study of asthma and allergy in childhood (ISAAC). It contains 11 questions pertaining to the diagnosis and risk factors of asthma. This questionnaire was adapted from ISAAC phase 1 and phase 2 with slight modification (questions indicating severity of asthma were excluded) and were

printed in both English and Marathi. The Marathi questionnaire was a pre validated translation. Questions were answered as either Yes or No. After permission from school authority, and clearance from local ethics committee, questionnaires were distributed to 1718 children. Explanation of each question was provided in the form. Questionnaire was to be answered by parents. After 2 reminders, 815 children returned the form (47.43%). Questionnaires were analysed and 62 out of 815 who answered yes for any question in section B were shortlisted as probable asthmatic and were evaluated at a tertiary teaching hospital in Pune. Detailed history and examination, PEFR was done. Out of these 62 children, 50 children were found to be asthmatic and they were included in the study and analysed. Asthma in this study population was defined as per the ISAAC study criteria that is - Presence of wheezing or whistling in the chest, chest sounded wheezy during exercise, and dry cough at night; apart from cough associated with cold or chest infection during past 12 months. Or if they answered yes to the question - has your child ever had asthma. Data was compiled and analysed using Statistical Package for Social Sciences (SPSS) software. Chi square, p test were used for determining statistical significance.

RESULTS

In this study 815 children were studied. Male to female ratio was 1.38 (male-57.9%, female -42.1%). Age wise distribution showed that 28.22% of children were in 5-8yrs age group, 33.12 % children in 9-11yrs age group and 38.65% children in 12-15 yrs. age group. 368 children were from Marathi medium (45.15 %) and 447 were from English medium schools (54.85 %).(Table-1).Overall Prevalence of asthma in our study was 6.13 %. Prevalence of asthma in age group of 5-8yrs (6.96%) was more than that in age group 9-11 yrs. (6.29%) and in age group 12-15 yrs. (5.39%). This difference was not statistically significant. In this study, overall male predominance was observed in the children with asthma. Prevalence of asthma in male children was 6.36% and in female children was 5.83 %. In age group 5-8 yrs. 7.6% female had asthma as compared to 6.52% males. In age

***Corresponding Author Dr. Manojkumar G. Patil**

Associate Professor, Dr D.Y. Patil Medical College, Hospital and Research Center& Dr. D.Y.Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.

group 9-11 yrs. and 12-15 yrs. male predominance was observed (6.53% vs 5.83% and 6.0% vs 5.83%) respectively (table-2). Out of 62 children labelled as probable asthmatics, 50 children were diagnosed as asthmatic after clinical evaluation. Out of 50 children, 30 were male and 20 were female. Family history of asthma and allergic diseases was present in 30 (60%), either asthma or allergic diseases in 17 (34%), and no family history of either in 3 (6%). Family history of asthma is an important predisposing factor in asthmatic children. In our study, 11.05% (90 out of 815) of cohort had family history of asthma, while 62% (31 out of 50) of asthmatic children had positive family history. This difference was statistically highly significant ($X^2 = 175.9, p < 0.0001$) (Table-3). More number of children with decrease PEFR had family history of asthma. 58.06% (18/31) of children with family history of asthma had decreased PEFR as compared to 52.6% (10/19) children without family history of asthma. This difference was statistically not significant ($p: 0.70$). (Table-4). In children with decreased PEFR it was found that 64.3% had positive family history of Asthma, 35.7% had no such history. Family history of other allergic diseases besides asthma also has significant association with prevalence of asthma. In this study 7.36% (60/815) of children had positive family history of allergic diseases. 56% (28/50) of asthmatics had family history of allergic disease other than asthma. This was highly significant ($p < 0.005$). (table-3) In children with family history of allergic diseases 57.1% (16/28) had decreased PEFR. While in children with no family history of allergic disease 54.5% (12/22) had decreased PEFR. ($P: 0.85$). Of asthmatic children 58% had history of atopic diseases. 56% of these had decrease in PEFR. In this study 8.05% (36 out of 447) of English medium students suffered from asthma as compared to 3.8% (14 out of 368) of students studying in Marathi medium. This was significant ($p < 0.05$) (table-1). Decrease in PEFR was seen in 58.3% of English medium students as compared to 50% in Marathi medium students ($p > 0.05$).

DISCUSSION

Asthma is chronic childhood disease with increasing prevalence over past decades. Environmental factors and genetic factors are contributing for the same. Pune is one of the most rapidly growing cities of India. In Pune, prevalence reported in 2004 by ISAAC phase 3 was 2.9%. While a study by Maria Cheraghi et. al. in 2009 and 2012 showed a prevalence of 5.4% and 6.7% respectively. (4). There was an increase in the prevalence of asthma by 130% in a decade. These were questionnaire based studies, where diagnosis of probable asthma was done by analysis of response sent by parents. The diagnosis was not confirmed by clinical examination. Prevalence of Asthma in present study was 6.13%. In our study, prevalence of asthma in age group 5-8 yrs. was 6.96%, in age group 9-11 yrs. was 6.29% and in age group 12-15 yrs. was 5.39%. A similar age wise prevalence of asthma of 7% among 6-7 yrs. old and 6.3% amongst 13-14 yrs. olds was reported from Pune. (4). An Iranian study by HadiBazzazi et al found prevalence of 7% in children aged 12-13 yrs. (6). As compared to 2012 Pune study, there was no increase in prevalence of asthma in our study. Unlike previous study, we had taken a detailed history and done examination and investigations of 62 probable asthmatics diagnosed by the questionnaire. We excluded 12 probable asthmatics as on examination we could not confirm the diagnosis of asthma in them. In spite of excluding 12 probable asthmatics from 62 probable asthmatics the prevalence of asthma in this study was 6.13%. We found male predominance in asthmatics. Prevalence of asthma in males was 6.36% and females were 5.83%. Male to female ratio was 1:1.5. Most of the studies have reported male predominance in prevalence of asthma. (7,8,9). Jain et. al. attributed it to increase bronchial lability in males. (10). S. K. Chhabra reported prevalence of 12% in males as compared to 11.2% in females. (8). Similarly Renata Gontijo et. al. reported prevalence of 8.1% and 6.1% respectively in males and females. (11). In our study 11.5% of children in cohort of 815 had family history of asthma. While in asthmatic group 62% children had family history of asthma. This correlation was highly significant ($p < 0.0001$). Similar strong relationship between family history of asthma and asthma has been reported previously. (8,10). First comprehensive study for inheritance of asthma was undertaken by Cook and Vander Veer in

1916. They came to conclusion that familial association was due to genetic component. (7,12). In this study we found strong association between family history of allergic diseases and asthma. 7.36% of 815 children had family history of allergic disorder, while 56% of asthmatic had family history of allergic diseases. Anzhela V. Glushkova et al. found that family history of allergic disease had highly significant association with prevalence of asthma in children. (13). HadiBazzazi et. al. that 30.2% of asthmatic children had family history of allergic disease with strong association between family history of allergic disease and asthma. (6) PEFR was performed at the time of confirmation of diagnosis. This was a baseline PEFR irrespective of the clinical symptoms. Hence the difference in PEFR value was not significant in children with predisposing factor and children without them. Out of 28 children with family history of allergic disease 57.1% had decreased PEFR as compared to 42.9% of children with no family history of allergic disease. 58.06% of children with family history of asthma had decrease in PEFR as compared to 41.9% of children without family history of asthma. We found that children who had both family history of asthma and family history of allergic disease had higher incidence of low PEFR as compared to those who had either or none. This may indicate higher chance of asthma being triggered by minor triggers in these children. Prevalence of asthma was 8.05% in students of English medium school, and was 3.8% in students of Marathi medium school.

Children from Marathi medium were from lower socioeconomic class as they were from government school and were from an area away from the city. Students from English medium were from a private school located in the crowded area of city. Dr Strachan's theory that children in larger families were exposed to more infection, resulting in healthier immune system that were less likely to mistake harmless substances for allergen. A similar difference of 7.3% and 5.8% in private and public schools respectively was reported by Maria Cheraghi et. al. (4). They stated that this was because students of private schools were from higher socioeconomic class as compared to public schools catering to lower socioeconomic class. The strength of this study was that after screening using ISAAC, probable asthmatics were examined and investigated. Children not confirming to diagnosis of asthma were not included. PEFR was performed in all asthmatic children, which was not the feature of other studies. Doing study in two schools was the limitation. We feel that more schools should be screened in this manner to get better representation of prevalence in the city. Such studies should be done at regular intervals so that trends in burden of disease can be determined and preventive methods be implemented. Future studies correlating air quality and life style with asthma is required for prevention planning.

Table No.1- Demographic Data of Study Population

| | | Cohort | Asthma | p value |
|---------------------|---------------|--------------|----------|---------|
| | | N=815 | N=50 | |
| Age (years) | 5-8 | 230 (28.22%) | 16 (32%) | 0.7483 |
| | 9-11 | 270 (33.12%) | 17 (34%) | |
| | 12-15 | 315 (38.65%) | 17 (34%) | |
| Sex | Male | 472 (57.91%) | 30 (60%) | 0.7578 |
| | Female | 343 (42.08%) | 20 (40%) | |
| | Male : Female | 1.38 : 1 | 1.5 : 1 | |
| Medium of education | Marathi | 368 (45.15%) | 14 (28%) | 0.01188 |
| | English | 447 (54.85%) | 36 (72%) | |

Table No.2 Prevalence of Asthma

| Age (years) | Male | | | p value | Female | | | p value | Total | | |
|-------------|-------|-----------|------------|---------|--------|-----------|------------|---------|-------|-----------|------------|
| | Total | Asthmatic | Prevalence | | Total | Asthmatic | Prevalence | | Total | Asthmatic | Prevalence |
| 5-8 | 138 | 9 | 6.52% | 0.9810 | 92 | 7 | 7.60% | 0.6122 | 230 | 16 | 6.96% |
| 9-11 | 153 | 10 | 6.53% | | 117 | 7 | 5.98% | | 270 | 17 | 6.29% |

| | | | | | | | | | |
|-------|-----|----|-------|-----|----|-------|-----|----|-------|
| 12-15 | 181 | 11 | 6.08% | 134 | 6 | 4.48% | 315 | 17 | 5.39% |
| Total | 472 | 30 | 6.36% | 343 | 20 | 5.83% | 815 | 50 | 6.13% |

Table No.3- Risk Factors for Asthma

| | Cohort N=815 | Asthma N=50 | P value |
|-------------------------------------|-----------------|----------------|----------|
| Family history of asthma | 90 (11.05%) | 31 (62%) | P<0.0001 |
| Family history of allergic disorder | 60 (7.36%) | 28 (56%) | P<0.0001 |

Table No.4 – Association of Risk Factors for Asthma& PEFR

| | LOW PEFR | NORMAL PEFR | P value |
|---------------------------------------|---------------|----------------|---------|
| | 28 | 22 | |
| Family history of asthma | 18 (64.3%) | 13 (59.1%) | 0.7072 |
| No Family history of asthma | 10 (35.7%) | 9 (40.9%) | |
| Family history of allergic disease | 16 (57.1%) | 12 (54.5%) | 0.8543 |
| No Family history of allergic disease | 12 (42.9%) | 10 (45.5%) | |
| Medium of Education: English | 21 (75%) | 15 (68.18%) | 0.5940 |
| Medium of Education: Marathi | 7 (25%) | 7 (31.81%) | |

REFERENCES

1. Morbidity and mortality report NCHS 2014- Health United States - Tables - 2014 Complete List. [online] Available at: <https://www.cdc.gov/nchs/hus/contents2014.htm>
2. Liu A, Covar R, Spahn J, Sicherer S. Childhood Asthma. In: Kliegman, RBehrman R, ed. by. Nelson TEXTBOOK of PEDIATRICS. 20th ed. Philadelphia, PA: Elsevier; 2016. p. 1095-1096.
3. National Asthma Education Program, (1991). Guidelines for the diagnosis management of asthma " Patient education. *Patient Education and Counseling*, [online] 18(1), pp.51-66. Available at: [http://dx.doi.org/10.1016/0738-3991\(91\)90079-k](http://dx.doi.org/10.1016/0738-3991(91)90079-k)
4. Cheraghi, M., Dadgarinejad, A. and Salvi, S. (2012). A Cross-Sectional Study to Find Prevalence and Risk Factors for Childhood Asthma in Pune City, India. *ISRN Public Health*, [online] 2012, pp.1-8. Available at: <http://dx.doi.org/10.5402/2012/361456>
5. Beasley, R., Ellwood, P. and Asher, I. (2003). International patterns of the prevalence of pediatric asthma. *Pediatric Clinics of North America*, [online] 50(3), pp.539-553. Available at: [http://dx.doi.org/10.1016/s0031-3955\(03\)00050-6](http://dx.doi.org/10.1016/s0031-3955(03)00050-6)
6. HadiBazzazi H, Gharagozlou M, Kassaiee M, Parsikia A, Zahmatkesh H. The prevalence of asthma and allergic disorders among school children in Gorgan. *Journal of Research in Medical Sciences* Jan & Feb 2007 [Internet]. 2007;12(1): 28-33. Available from: <http://jrms.mui.ac.ir/index.php/jrms/article/view/365/285>
7. Qureshi, U., Qureshi, U., Bilques, S., ulHaq, I., Khan, M. and Qurieshi, M. (2016). Epidemiology of bronchial asthma in school children (10-16 years) in Srinagar. *Lung India*, [online] 33(2), p.167. Available at: <http://dx.doi.org/10.4103/0970-2113.177442>
8. Chhabra, S., Gupta, C., Chhabra, P. and Pal, S. (1998). Prevalence of Bronchial Asthma in Schoolchildren in Delhi. *Journal of Asthma*, [online] 35(3), pp.291-296. Available at: <http://dx.doi.org/10.3109/02770909809068220>
9. Mvula, M., Larzelere, M., Kraus, M., Moisiejewicz, K., Morgan, C., Pierce, S., Post, R., Nash, T. and Moore, C. (2005). Prevalence of Asthma and Asthma-Like Symptoms in Inner-City Schoolchildren. *Journal of Asthma*, [online] 42(1), pp.9-16. Available at: <http://dx.doi.org/10.1081/jas-200044746>
10. Jain, A., Vinod Bhat, H. and Acharya, D. (2010). Prevalence of

bronchial asthma in rural Indian children: A cross sectional study from South India. *The Indian Journal of Pediatrics*, [online] 77(1), pp.31-35. Available at: <http://dx.doi.org/10.1007/s12098-009-0308-6>

11. Lima, R., Pastorino, A., Casagrande, R., Sole, D., Leone, C. and Jacob, C. (2007). Prevalence of asthma, rhinitis and eczema in 6-7 years old students from the western districts of SÃo Paulo City, using the standardized questionnaire of the "International Study of Asthma and Allergies in Childhood" (ISAAC)-phase IIIB. *Clinics*, [online] 62(3). Available at: <http://dx.doi.org/10.1590/s1807-59322007000300005>
12. Cooke, R. and Vander veer, A. (1916). Human sensitisation. *J. Immunol*, [online] 1(3), pp.201-305. Available at: <http://www.jimmunol.org/content/1/3/201>.
13. Glushkova, A. and Grjibovski, A. (2008). Prevalence and Correlates of Asthma among Children in Central St. Petersburg, Russia: Cross-sectional Study. *Croatian Medical Journal*, [online] 49(6), pp.741-750. Available at: <http://dx.doi.org/10.3325/cmj.2008.49.741>